



Polycystic Horseshoe Kidney and Polycystic Liver Disease

Gürsel Yıldız¹, Ender Hür², Kemal Mağden², Sumru Tanju Şen³, Abdurkerim Yılmaz⁴ and Alp Gürkan^{5*}

¹Department of Nephrology, Istanbul Okan University School of Medicine, Turkey

²Department of Nephrology, Bulent Ecevit University School of Medicine, Turkey

³Department of Radiology, Ataturk State Hospital, Turkey

⁴Department of Gastroenterology, Cumhuriyet University School of Medicine, Turkey

⁵Department of General Surgery, Istanbul Okan University School of Medicine, Turkey

Abstract

Horseshoe kidney is a renal fusion anomaly during embryogenesis with incidence ranges of 1 in 400 to 1 in 1800 live births. Adult polycystic kidney disease is a hereditary disorder and transmitted in autosomal dominant pattern with a reported incidence of 1 in 1000 to 1 in 5000 cases. Polycystic horseshoe kidney is a very rare occurrence with incidence ranges of 1 in 134 000 to 1 in 8 000 000 live births. We here present a patient who was admitted at the hospital with headache, hypertension and polycystic horseshoe kidney.

Keywords: Horseshoe kidney; Polycystic kidney disease; Polycystic liver disease

Introduction

Horseshoe kidney is a fusion anomaly occurring during embryogenesis in which the kidneys are conjuncted in the lower poles of the median line with an incidence between 1/400 and 1800 [1]. Adult type autosomal Dominant Polycystic Kidney Disease (ADPKD) is a hereditary disorder developing due to polycystin 1 (ADPKD1) and polycystin 2 (ADPKD2) with a reported incidence of 1/1000-5000 [2]. Polycystic horseshoe kidney anomaly is a very rare disorder with an incidence between 1/134 000 and 1/8 000 000 [3]. Polycystic horseshoe kidney is considered as two separate diseases.

OPEN ACCESS

*Correspondence:

Alp Gürkan, Department of General Surgery, Istanbul Okan University School of Medicine, İçmeler Mahallesi, Aydınlı Yolu Cad, Aydemir Sk. No: 2, 34947 İçmeler-Tuzla, Istanbul, Turkey, Tel: 0090 216 444 9863; E-mail: algurkan@okan.edu.tr

Received Date: 01 Feb 2019

Accepted Date: 01 Apr 2019

Published Date: 04 Apr 2019

Citation:

Yıldız G, Hür E, Mağden K, Tanju Şen S, Yılmaz A, Gürkan A. Polycystic Horseshoe Kidney and Polycystic Liver Disease. *World J Surg Surgical Res.* 2019; 2: 1115.

Copyright © 2019 Alp Gürkan. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Case Presentation

A 54 years old female patient presented to our hospital with complaint of headache. From her history, we learnt that she had been treated for hypertension for 2 years. We found that our patient was going to the controls irregularly and used angiotensin receptor blocker + hydrochlorothiazide group antihypertensive medication. On her familial history, there was history of hypertension in her mother and there was not a history of kidney disease in her family. Her blood pressure was 180/100 mmHg without any additional pathology. On the laboratory examinations, creatinine value was found as 0.8 mg/dL. Full blood count and the other biochemical serum values were in the normal ranges. Microscopic hematuria was found in her urine examination. On the abdominal ultrasonography, it was found that the size of both kidneys increased, there were numerous numbers of cysts in the liver and kidneys and lower poles of the kidneys were conjuncted. Furthermore, the patient underwent computed abdominal tomography. On her computed tomography; enlarged kidneys, numerous kidney cysts with various sizes and horseshoe kidney anomalies (Figure 1) and multiple liver cysts (Figure 2) were reported. Her mother and siblings were screened with kidney ultrasonography for ADPKD, but no polycystic kidney disease was found.

Discussion

Horseshoe kidney, which occurs in the lower poles of the kidneys during embryogenesis, is probably the most common form of the renal fusion anomalies. Its incidence has increased in the cases with Turner's syndrome and in men [4]. Although, horseshoe kidney cases are asymptomatic, yet they can be seen with stone disease, ureteropelvic junction syndrome and renovascular hypertension [5-8].

Adult polycystic kidney disease is characterized with diffuse cystic lesions in the kidneys and other viscera. Mutation in the PKD1 gene which is located in the 16th chromosome and PKD2




Figure 1: Multiple cysts in various sizes and horseshoe anomaly in both the kidneys are seen on computed abdominal tomography.




Figure 2: Multiple cysts in various sizes in the liver are seen on computed abdominal tomography (polycystic liver).

gene which is located in the 4th chromosome, coding polycystin-1 and polycystin-2 in the kidneys and other tissues play a role in the pathogenesis [9-11]. Since PKD1 and PKD2 genes may be found also in the non-renal tissues, polycystic kidney disease must be considered as a systemic disorder. In adult polycystic disease, systemic involvement may be seen such as diffuse cystic lesions in the liver, colonic diverticula, valvular heart disorders and cerebral artery aneurysms [12-15]. There were diffuse hepatic cysts in our patient. Adult polycystic kidney disease may affect all the ethnic groups of ADPKD cases, 5% are seen without a familial history [3]. This probably reflects the new mutation causing to a new cohort likewise our case. So far, no any genetic association was described between PKD gene loci and horseshoe kidney anomaly.

We could not find data in the literature about clinical progression of polycystic horseshoe kidney. Diagnosis for polycystic horseshoe kidney is often established incidentally with ultrasonography during the routine screening or screening of the persons having familial ADPKD history.

Polycystic horseshoe kidney is, extremely rare with about 20 cases reported in the literature. Polycystic horseshoe kidney presented by findings of the combination of two distinct renal disorders, loss of kidney function and end stage renal failure [16]. Occasionally polycystic horseshoe kidney presented by such as subjective findings increased belly volume, abdominal pain, headache and hematuria. In the patients with polycystic horseshoe kidney disease, treatment is in the form of individual therapy especially for hypertension and serious complications including pain, hematuria, infection, nephrolithiasis, diverticulosis, and arterial aneurysms. Isthmus undivided bilateral nephrectomy has been reported in polycystic horseshoe kidney patients [16].

References

1. Wilson C, Azmy AF. Horseshoe kidney in children. *Br J Urol.* 1986;58(4):361-3.

2. Çağlar K, Kibar Y, Tahmaz L, Tekin A, Yenicesu M, Vural A. Polycystic horseshoe kidney. *Clin Nephrol.* 2001;55(6):487-8.
3. Brumm FA, Becker M, Uglione A, Da Ros CT. Polycystic horseshoe kidney. *J Urol.* 1997;158(6):2229.
4. Kleta R, Brämswig JH. Horseshoe kidney and Turner syndrome. *Nephrol Dial Transplant.* 2000;15(7):1094.
5. Fekak H, Mezzour MH, Rabii R, Joual A, Bennani S, El Mrini M. Management of horseshoe kidney based on a series of 36 cases. *Prog Urol.* 2004;14(4):485-8.
6. Bozic B, Persec Z, Persec J, Borkovic Z, Galesic K. Diagnostic and treatment of renovascular hypertension in the patient with horseshoe kidney. *Acta Med Croatica.* 2005;59(2):137-42.
7. Işksalan Özbülül N, Parlak S. A Case of Horseshoe Kidney with Ureteropelvic Junction Obstruction: Diagnosis by Multi-Detector Row Ct Urography. *Firat Med J.* 2009;14(1):72-4.
8. Kırkpankur A, Altun B, Yılmaz Mİ, Yenice M. A Hypertensive Patient With Polycystic Horseshoe Kidney. *Turk Neph Dial Transpl.* 2009;18(2):87-9.
9. The International Polycystic Kidney Disease Consortium: Polycystic Kidney Disease. The complete structure of the PKD1 and its protein. *Cell.* 1995;81(2):289-98.
10. Peters DJ, Spruit L, Saris JJ, Ravine D, Sandkuijl LA, Fossdal R, et al. Chromosome 4 localization of a second gene for autosomal dominant polycystic kidney disease. *Nature Genet.* 1993;5(4):359-62.
11. Kimberling WJ, Kumar S, Gabow PA, Kenyon JB, Connolly CJ, Somlo S. Autosomal dominant polycystic kidney disease: localization of the second gene to chromosome 4q13-q23. *Genomics.* 1993;18(3):467-72.
12. Hoevenaren IA, Wester R, Schrier RW, McFann K, Doctor RB, Drenth JP, et al. Polycystic liver: clinical characteristics of patients with isolated polycystic liver disease compared with patients with polycystic liver and autosomal dominant polycystic kidney disease. *Liver Int.* 2008;28(2):264-70.
13. Peña JM, Pernaute R, Vicente de Vera C. Is ADPKD associated with small-

- bowel diverticular disease? *Nephrol Dial Transplant*. 2000;15(11):1890-1.
14. Lumiaho A, Ikäheimo R, Miettinen R, Niemitukia L, Laitinen T, Rantala A, et al. Mitral valve prolapse and mitral regurgitation are common in patients with polycystic kidney disease type 1. *Am J Kidney Dis*. 2001;38(6):1208-16.
15. Pirson Y, Chauveau D, Torres V. Management of cerebral aneurysms in autosomal dominant polycystic kidney disease. *J Am Soc Nephrol*. 2002;13(1):269-76.
16. Yao XL, Jin BY, Wang YM. Isthmus Undivided Bilateral Nephrectomy in a Patient with Polycystic Horseshoe Kidney. *Chin Med J (Engl)*. 2015;128(5):698-9.