



## A Rare Case of Intestinal Obstruction Transomental Hernia

Mouhafid FE\*, Abdoulmalik S, Ramraoui MES, Fihri JF, Baba H, Lahkim M, Ahmed ELK and Rachid E

Department of General Surgery, Avicenne Military Teaching Hospital, Morocco

### Abstract

We report the case of a 70-year-old patient with a transomental hernia complicated by intestinal obstruction who underwent median laparotomy.

**Key words:** Internal hernia; Transomental hernia; Intestinal obstruction; Small intestine

### Introduction

Internal hernias are defined as a protrusion of hollow viscera through an intraperitoneal orifice and which remains in the abdominal cavity [1,2]. They constitute a rare cause of intestinal obstruction. In fact, they can be responsible for 0.2% to 5.8% of small intestinal obstructions [3]. Depending on their physio pathological mechanism, we distinguish several types, among which the transomental hernia constitutes one of the most exceptional and represents only 2% of internal hernias [1]. The trans-omental hernia is said to be true if the incarceration of the loops takes place in an orifice located near the free edge of the greater omentum on the right side [1]. The etiological diagnosis of a transomental hernia is not easy. The diagnosis is most often made intraoperatively in a patient with intestinal obstruction.

### Clinical Case

It was a 70-year-old man who was seen in the emergency room for abdominal pain with vomiting and cessation of materials and gas. During questioning, the patient reported that the symptoms began approximately 24 hours ago, with a twisting pain located in the periumbilical area radiating to the flank with food and then bilious vomiting, all evolving in an afebrile context. He would be type 2 diabetic under treatment.

### OPEN ACCESS

#### \*Correspondence:

Mouhafid FE, Department of General Surgery, Avicenne Military Teaching Hospital, Marrakech, Morocco,

Received Date: 05 Nov 2024

Accepted Date: 11 Nov 2024

Published Date: 04 Dec 2024

#### Citation:

Mouhafid FE, Abdoulmalik S, Ramraoui MES, Fihri JF, Baba H, Lahkim M, et al. A Rare Case of Intestinal Obstruction Transomental Hernia. *World J Surg Surgical Res.* 2024; 7: 1578.

**Copyright** © 2024 Mouhafid FE. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### The patient's clinical examination revealed

On general examination: general condition WHO stage 2, Glasgow at 15, vital constants T: 37.9; CF: 94; BP: 110/64.

### On physical examination

a distended abdomen, tympanic, diffusely painful on palpation of the free parietal hernial orifices and rectal examination with an empty rectal ampulla.

### The paraclinical examination

On the biological assessment, an elevation of the CRP to 160, hyperleucocytosis with a neutrophil predominance, no ionic disorders.

At the ASP we had hail-like hydro-aerial levels, we completed with Abdominal CT which showed hydro-aerial levels with small bowel distension with disparity in caliber without sign of damage to the loops.

The patient underwent a median laparotomy above and below the umbilical with the intra-op observation of an omental hernia by incarceration of ileal loops without signs of necrosis through a defect in the greater omentum of approximately 4 cm without other anomalies to exploration (Figure 1). We therefore carried out extrication of the ileal loops with closure of the defect using separate points with 3/0 vycril (Figure 2).

The postoperative course was simple and the patient was allowed to leave the hospital on the 7th post-operative day.



**Figure 1:** The surgical pictures of Transomental internal hernias.



**Figure 2:** Viable small intestine.

## Discussion

Internal hernias constitute an unusual pathology in current practice; among its different varieties, transomental hernias are one of the rarest and represent only 2% of cases of internal hernias [1].

Transomental internal hernias are observed in patients who have no previous abdominal surgery, as is the case in our patient. They occur most often in patients aged over 50 [4] as in our case. These hernias can be seen in both sexes but are more common in males.

In our case, the hernia was revealed by an occlusive syndrome which is the most common mode of manifestation of this type of hernia when the incarcerated small bowel loops are strangulated [5]. The previously difficult preoperative etiological diagnosis is made easier by the more frequent use of CT. MRI is a better diagnostic tool than CT but is most often not available in an emergency. In our case, the etiological diagnosis of the occlusion was made intraoperatively.

The standard treatment for this type of pathology consists of a laparotomy to remove the obstacle and closure of the defect to avoid the occurrence of intestinal necrosis or recurrence, hence the need for early treatment.

## Conclusion

Transomental hernias constitute a rare pathology in current practice whose etiological diagnosis is facilitated by imaging techniques. They can constitute a medico-surgical emergency in the event of a diagnostic and therapeutic delay.

## References

1. Inoue Y, Nakamura H, Mizumoto S, Akashi H. Leser sac hernia through the gastrocolic ligament. CT diagnosis. *Abdom imaging*. 1996;21(2):145-7.
2. Mathias J, Phi I, Bruot O, Ganne PA, Laurent V, Regent D. Hernies internes. EMC (Elsevier Masson SAS), Paris, Radiodiagnostic – Appareil digestif. 2008;33-15-A-37.
3. Takeyama N, Gokan T, Ohgiya Y, Satoh S, Hashizune T, Hataya K, et al. CT of internal hernias. *Radiographics*. 2005;25(4):997-1015.
4. Watt PC. Transomental hernia causing intestinal obstruction in an elderly patient. *Postgrad Med J* 1983;59(698):790.
5. Zissin R, Hertz M, Gayer G, Paran H, Osadchy A. Internal hernia as a cause of small bowel obstruction: CT findings in 11 adult patients. *Br J Radiol*. 2005;78(933):796-802.