



Primary Squamous Cell Carcinoma of the Sigmoid Colon Revealed by Acute Intestinal Obstruction: A Case Report and Literature Review

Ahmed Zeroual*, Mohammed Essaid Ramraoui, Ahmed EL Guazzar, Mohamed Jawad Fassi EL Fihri, Hicham Baba, Mohammed Lahkim, Ahmed El Khader and Rachid El Barni

Department of Visceral Surgery, Avicenne Military Hospital, Marrakech, Morocco

Abstract

Primary squamous cell carcinoma (SCC) of the colon is an exceptional tumor. Fewer than 150 cases have been reported in the literature up to 2014. Besides its rarity, it is often associated with other digestive neoplasms. We report the case of a 54-year-old patient with primary SCC of the colon. This case allows us to discuss the anatomic-clinical and therapeutic characteristics, as well as the etiopathogenic hypotheses of this uncommon entity.

Keywords: Colon; Squamous cell carcinoma; Prognosis

Introduction

Colonic localization of primary squamous cell carcinoma (SCC) is extremely rare. The histogenesis and evolutionary potential of such a tumor remain unresolved [1-3].

The purpose of this work is to highlight that, despite its exceptional topographical occurrence, this diagnosis should not be overlooked. Therapeutic choices and comprehensive management can only be considered within a multidisciplinary framework. We also aim to report this case and explore similar cases to compare them with existing literature.

OPEN ACCESS

*Correspondence:

Ahmed Zeroual, Department of Visceral Surgery, Avicenne Military Hospital, Marrakech, Morocco,
E-mail: drzeroualahmed1010@gmail.com

Received Date: 31 Jan 2025

Accepted Date: 14 Feb 2025

Published Date: 17 Feb 2025

Citation:

Zeroual A, Ramraoui ME, Guazzar AEL, Fassi EL Fihri MJ, Baba H, Lahkim M, et al. Primary Squamous Cell Carcinoma of the Sigmoid Colon Revealed by Acute Intestinal Obstruction: A Case Report and Literature Review. *World J Surg Surgical Res.* 2025; 8: 1588.

Copyright © 2025 Ahmed Zeroual. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Case Presentation

A 64-year-old male, with no notable medical history, initially presented with abdominal pain associated with constipation and unquantified weight loss. The patient was admitted to the emergency department for acute intestinal obstruction. Rectal examination revealed an empty rectal ampulla without signs of bleeding.

Abdominal CT scans in axial and coronal sections revealed irregular, stenosing wall thickening of the sigmoid colon, enhanced by contrast medium, indicating a tumor causing colonic obstruction upstream, without evidence of locoregional extension or hepatic metastases.

The patient underwent elective colostomy for decompression to allow for bowel preparation and histopathological diagnosis.

Histopathological analysis of the biopsy revealed a poorly differentiated carcinomatous process. A left colectomy was performed for oncological purposes with colorectal anastomosis following staging and a multidisciplinary team discussion.

Macroscopically, the surgical specimen showed a 3.4 cm × 4.6 cm exophytic colonic tumor with a whitish-gray appearance on cross-section. Extensive tumor sampling revealed a carcinomatous proliferation consisting of nests and cords, without glandular structures.

The tumor cells were large, with intercellular bridges and keratinization, eosinophilic cytoplasm, and nuclei showing moderate to marked atypia with mitotic figures and several keratin pearls. Vascular emboli were present, but perineural invasion was absent. Lateral and circumferential margins were clear. Lymph node dissection revealed metastasis with one capsular breach (2/28 nodes positive). The tumor was classified as pT3N1bM0.

Immunohistochemical analysis concluded that the tumor was a well-differentiated, infiltrating, keratinizing SCC of the colon.




Figure 1: Axial and coronal scans revealed an irregular, stenosing thickening of the sigmoid wall, enhanced after contrast injection, of tumoral origin, causing an upstream colonic obstruction without signs of locoregional extension or hepatic metastases.




Figure 2: Histological section showing a poorly differentiated carcinomatous proliferation (H&E x200).

Clinical examination of skin and mucosal surfaces, along with a comprehensive staging workup (thoracoabdominal-pelvic CT scan and brain MRI), showed no secondary localization, confirming the primary nature of the colonic SCC.

The patient underwent adjuvant therapy, with favorable postoperative evolution. The patient remains under follow-up in medical oncology six months post-treatment.

Discussion

Primary SCC of the colon is extremely rare, accounting for less than 0.5% of all colorectal tumors, with an estimated incidence of 0.1% [1,2]. The etiopathogenesis of colonic SCC remains unclear. It may arise from a multipotent stem cell or develop from squamous metaplasia secondary to chronic irritation [2-4]. Supporting this hypothesis is the frequent association of colonic SCC with chronic inflammatory conditions, particularly ulcerative colitis, where the relative incidence is 1.7% compared to 0.25% to 0.5% in the general population [5]. Other potential causes include parasitic colitis, such as amebiasis or schistosomiasis, and chronic colocutaneous fistulas, which may lead to squamous metaplasia [5,6].

Colonic SCC typically occurs around the fifth decade of life with a male predominance (sex ratio: 2) [6,7]. It preferentially affects the cecum and right colon [7], as in our case, and less frequently the left colon. Clinically, symptoms are similar to those of adenocarcinomas [2,4], with a diagnostic delay ranging from six weeks to 12 months [6]. In some cases, metastases—often hepatic or pulmonary—are the first manifestation [7,8].

The diagnostic criteria for colonic SCC are stringent. The tumor must be located at least 7 cm above the anoperineal line to exclude SCC of the anal canal and lower rectum. Histologically,




Figure 3: Strong expression of tumor cells with the P40 antibody (IHC x200).

adenosquamous carcinoma—a rare entity combining glandular and squamous components—must also be ruled out. This requires extensive tumor sampling. Before concluding the primary nature of the tumor, a search for SCC in other organs must exclude metastatic origins [1,2,8,9].

In 10% of cases, colonic SCC is associated with another digestive tumor, often a synchronous adenocarcinoma [9,10]. Associations with extra digestive cancers, including ovarian, endometrial, prostatic, and breast carcinomas, have also been reported [10,11].

Adjuvant treatment results are inconsistent. The efficacy of radiotherapy, alone or combined with chemotherapy, remains unestablished due to the tumor's rarity [1,2,10]. Colonic SCC has a worse prognosis compared to glandular carcinoma. Mortality occurs within the first year in 52% of cases [10]. Factors associated with poor outcomes include left-sided tumor location, ulcerated lesions, nodal metastases, poor differentiation, and TNM stage IV [1,10].

Conclusion

Primary colonic SCC is an exceptional tumor with unclear etiopathogenesis. Its carcinogenesis remains poorly understood, and its prognosis is less favorable than that of typical adenocarcinomas.

Conflicts of Interest

The authors declare no conflicts of interest.

Author Contributions

All authors contributed to the article's preparation. All authors read and approved the final manuscript.

References

1. Santosh K. Mondal, Primary squamous cell carcinoma of the caecum: a case report. *J Cancer Res Ther.* 2011;5(4):5.
2. Michelassi F, Mishlove LA, Stipa F, Block GE. Squamous-cell carcinoma of the colon: experience at the University of Chicago, review of the literature, report of two cases. *Dis Colon Rectum.* 1988;31(3):228-35.
3. Frizelle FA, Hobday KS, Batts KP, Nelson H. Adenosquamous and squamous carcinoma of the colon and upper rectum: a clinical and histopathologic study. *Dis Colon Rectum.* 2001;44(3):341-6.
4. Daldoul S, Moussi A, Zaouche A. Carcinome épidermoïde primitif du colon: à propos d'un cas exceptionnel associé à un adénocarcinome gastrique. *J Afr Hépatol Gastroentérol.* 2011;5:221-3.
5. Pittella JE, Torres AV. Primary squamous-cell carcinoma of the cecum and ascending colon: report of a case and review of the literature. *Dis Colon Rectum.* 1982;25(5):483-7.
6. Lyttle JA. Primary squamous carcinoma of the proximal large bowel: report of a case and review of the literature. *Dis Colon Rectum.* 1983;26(4):279-

- 82.
7. Vezeridis MP, Herrera LO, Lopez GE, Ledesma EJ, Mittleman A. Squamous-cell carcinoma of the colon and rectum. *Dis Colon Rectum*. 1983;26(3):188-91.
 8. Chulia F, Camps C, Rodriguez A, Medina E, Tuset J. Epidermoid carcinoma of the colon: description of a lesion located in the hepatic flexure. *Dis Colon Rectum*. 1996;29(10):665-7.
 9. Michelassi F, Montag AG, Block GE. Primary squamous cell carcinoma in ulcerative colitis: report of a case. *Dis Colon Rectum*. 1988;31(4):323-6.
 10. Zoran J, Iva L, Monika U, Mario L, Božo K, Davor T. Primary acantholytic squamous cell carcinoma of the cecum: a case report. *Diagn Pathol*. 2011;6:5.
 11. Palvio DH, Sorensen FB, Kløve-Mogensen M. Stem cell carcinoma of the colon and rectum: report of two cases and review of the literature. *Dis Colon Rectum*. 1985;28(6):440-5.